



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 31/365</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 97/38689</b> <b>(43) International Publication Date:</b> 23 October 1997 (23.10.97)
<b>(21) International Application Number:</b> PCT/EP97/01800 <b>(22) International Filing Date:</b> 10 April 1997 (10.04.97)  <b>(30) Priority Data:</b> 9607564.3                      12 April 1996 (12.04.96)                      GB 9622028.0                      24 October 1996 (24.10.96)                      GB  <b>(71) Applicant (for all designated States except US):</b> NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> HAEBERLIN, Barbara [CH/CH]; Haselrain 77, CH-4125 Riehen (CH). MAK, Ching, Pong [GB/CH]; Ob dem Fichtenrain 20, CH-4106 Therwil (CH). MEINZER, Armin [DE/DE]; Fliederweg 2, D-79426 Buggingen (DE). VONDERSCHER, Jacky [FR/FR]; 29, rue de Bruebach, F-68400 Riedisheim (FR).  <b>(74) Agent:</b> ROTH, Bernhard, M.; Novartis AG, Lichtstrasse 35, CH-4002 Basel (CH).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> ENTERIC-COATED PHARMACEUTICAL COMPOSITIONS OF MYCOPHENOLATE  <b>(57) Abstract</b> <p>This invention provides a pharmaceutical composition comprising a mycophenolate salt, the composition being adapted to release mycophenolate in the upper part of the intestinal tract.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## ENTERIC-COATED PHARMACEUTICAL COMPOSITIONS OF MYCOPHENOLATE

This invention relates to mycophenolic acid.

5 Mycophenolic acid, also referred to herein as MPA, was first isolated in 1896, and has been extensively investigated as a pharmaceutical of potential commercial interest. It is known to have anti-tumor, anti-viral, immunosuppressive, anti-psoriatic, and anti-inflammatory activity [see e.g. W.A. Lee et al, *Pharmaceutical Research* (1990), 7, p. 161 - 166 and references cited therein]. Publications have appeared on MPA as an anti-cancer agent by Lilly scientists, see e.g. M.J. Sweeney et al., *Cancer Research* (1972), 32, 1795-1802, and  
10 by ICI scientists, see e.g. GB 1,157,099 and 1,203,328 and as an immunosuppressant agent see e.g. A. Mitsui et al. *J. Antibiotics* (1969) 22, p. 358-363. In the above-mentioned article by W.A. Lee et al it is stated that attempts have been made to increase the bio-availability or specificity of MPA by making derivatives. The poor bioavailability of the acid was thought to be caused by undetermined factors such as drug complexation in the gastro-  
15 intestinal lumen, a narrow absorption window, metabolism before absorption etc.. The preparation of the morpholinoethyl ester, also known as mycophenolate mofetil (sometimes referred to herein as MMF), was described which had considerably higher bioavailability than MPA (100% for MMF and 43% for MPA). This derivative has been recently introduced commercially as an immunosuppressant for the treatment or prevention of organ  
20 or tissue transplant rejection, at daily dosages of from about 200 mg to about 3 grams p.o., e.g. about 2 g p.o. Patient compliance with MMF is not ideal, inter alia, because of side-effects e.g. gastro-intestinal side effects, the origin of which is not known.

We have now found, after exhaustive testing, that mycophenolate salts when enteric coated or adapted to be released in the upper part of the intestines, e.g. in the duodenum, jejunum  
25 and/or ileum, are effective, well-tolerated, pharmaceuticals particularly for immunosuppressive indications especially for the treatment or prevention of organ, tissue

or cellular allograft or xenograft rejection, e.g. after transplant, or the treatment or prevention of immune-mediated diseases (autoimmune diseases) and have interesting bioavailability and stability characteristics. Moreover fewer unit dosage forms are required to be administered than for MMF, leading to easier administration.

5 The present invention provides in one aspect a pharmaceutical composition comprising a myophenolate salt, the composition being adapted to release mycophenolate in the upper part of the intestinal tract (hereinafter referred to as a composition of the invention). The composition may be adapted in any conventional manner, preferably with means adapted to prevent release of the mycophenolate in the stomach and to ensure release in the upper  
10 part of the intestinal tract. In a further aspect the invention provides a pharmaceutical composition comprising a coated pharmaceutically acceptable mycophenolate salt.

Such salts are cationic salts, e.g. of alkali metals, especially the sodium salts. Sodium mycophenolate salts are known, e.g. in South African Patent 68/4959. We prefer to use the mono-sodium salt. This may be obtained in crystalline form by recrystallization from  
15 acetone/ethanol if necessary with water; Mpt. 189 - 191°C.

The invention provides, more specifically, a solid enteric-coated composition in unit dose form for oral application, the core of the composition containing sodium mycophenolate in solid or liquid form.

The term "core" comprises sodium mycophenolate (or other cationic salt) if desired in  
20 admixture with further physiologically acceptable material, that can be surrounded by an enteric-coating. The term "core" comprises, in a wide sense, not only tablets, pellets or granules but also capsules, e.g. soft or hard capsules of gelatine or starch. Such cores may be produced in conventional manner. We have found that the mycophenolate salts, particularly the sodium salt, are particularly interesting for the production of tablets. When  
25 tablet cores are used they have preferably a hardness of from ca. 10 to 70 N.

The pellets or granules may, after application of the enteric-coating as described hereinafter may be used as such or to fill capsules, e.g. hard gelatine capsules. If desired the capsules may be alternatively enteric-coated, e.g. in conventional manner.

5 Other pharmaceutically acceptable ingredients may be present in the cores, e.g. those conventionally used in the preparation of pharmaceutically compositions, e.g. fillers, e.g. lactose, glidants, e.g. silica, and lubricants, e.g. magnesium stearate.

10 The term "enteric coating" comprises any pharmaceutically acceptable coating preventing the release of the active agent in the stomach and sufficiently disintegrating in the intestine tract (by contact with approximately neutral or alkaline intestine juices) to allow the resorption of the active agent through the walls of the intestinal tract. Various in vitro tests for determining whether or not a coating is classified as an enteric coating have been published in the pharmacopoeia of various countries.

15 More specifically, the term "enteric coating" as used herein refers to a coating which remains intact for at least 2 hours, in contact with artificial gastric juices such as HCl of pH 1 at 36 to 38°C and preferably thereafter disintegrates within 30 minutes in artificial intestinal juices such as a  $\text{KH}_2\text{PO}_4$  buffered solution of pH 6.8.

The thickness of the coating may vary and depends inter alia on its permeability in water and acids. A typical coating may be about 16-30, e.g. 16-20 or to 25, mg on a size 1 gelatine capsule. Similar thicknesses may be applied in other formulations.

20 In general satisfactory results are obtained with a coating of 5 - 100  $\mu\text{m}$ , preferably 20 - 80  $\mu\text{m}$  thickness. The coating is suitably selected from macromolecular polymers. Suitable polymers are listed in e.g. L. Lachman et al. The Theory and Practice of Industrial Pharmacy, 3rd Ed, 1986, p. 365 - 373, H. Sucker et al, Pharmazeutische Technologie, Thieme, 1991, p. 355 - 359, Hagers Handbuch der pharmazeutischen Praxis, 4th Ed. Vol.  
25 7, pages 739 to 742 and 766 to 778, (Springer Verlag, 1971) and Remington's Pharmaceutical Sciences, 13th Ed., pages 1689 to 1691 (Mack Publ., Co., 1970) and

comprise e.g. cellulose ester derivatives, cellulose ethers, acrylic resins, such as methylacrylate copolymers and copolymers of maleic acid and phthalic acid derivatives.

The preferred films are made from cellulose acetate phthalate and trimellitate; methacrylic acid copolymers, e.g. copolymers derived from methylacrylic acid and esters thereof, containing at least 40% methylacrylic acid; and especially hydroxypropyl methylcellulose phthalate.

Methylacrylates include those of molecular weight above 100,000 daltons based on, e.g. methylacrylate and methyl or ethyl methylacrylate in a ratio of about 1:1. Typical products include Endragit L, e.g. L 100-55, marketed by Rohm GmbH, Darmstadt, Germany.

Typical cellulose acetate phthalates have an acetyl content of 17-26% and a phthalate content of from 30-40% with a viscosity of ca. 45-90cP.

Typical cellulose acetate trimellititates have an acetyl content of 17-26%, a trimellityl content from 25 - 35 % with a viscosity of ca. 15-20 cS. An example of an appropriate cellulose acetate trimellitate is the marketed product CAT (Eastman Kodak Company, USA).

Hydroxypropyl methylcellulose phthalates, typically have a molecular weight of from 20,000 to 100,000 daltons e.g. 80,000 to 130,000 daltons, e.g. a hydroxypropyl content of from 5 to 10%, a methoxy content of from 18 to 24% and a phthalyl content from 21 to 35%.

An example of an appropriate cellulose acetate phthalate is the marketed product CAP (Eastman Kodak, Rochester N.Y., USA).

Examples of suitable hydroxypropyl methylcellulose phthalates are the marketed products having a hydroxypropyl content of from 6-10%, a methoxy content of from 20-24%, a phthalyl content of from 21-27%, a molecular weight of about 84,000 daltons known under the trade mark HP50 and available from Shin-Etsu Chemical Co. Ltd., Tokyo, Japan, and

having a hydroxypropyl content, a methoxy content, and a phthalyl content of 5-9%, 18-22% and 27-35% respectively, and a molecular weight of 78,000 daltons, known under the trademark HP55 and available from the same supplier.

A preferred coating is HP 50.

- 5        The enteric coating may be carried out in conventional manner, e.g. so that the cores are sprayed with a solution of the enteric-coating.

Suitable solvents for the enteric-coating are for example organic solvents, e.g. an alcohol such as ethanol, a ketone such as acetone, halogenated hydrocarbons such as  $\text{CH}_2\text{Cl}_2$  or mixtures of such solvents, e.g. ethanol /acetone, e.g. 1:1 to 10:1.

- 10       Conveniently a softener such as di-n-butylphthalate or triacetin is added to such a solution, e.g. in a ratio of coating material to softener of from 1: about 0.05 to about 0.3.

If desired for cellulose phthalates and other acidic coating materials an ammonium salt may be found and an aqueous solution may be used.

A fluidized bed coater may be used for coating.

- 15       Conveniently the cores are treated at room temperature or warmed up to 40°C e.g. by means of warm air of 40° up to 70°C, before spraying. To avoid a sticking of the cores the spray procedure is preferably interrupted at certain time intervals and the cores then warmed up again. It is, however, also possible to proceed without interruption of the spray procedure, e.g. by automatic regulation of the spray amount taking into account the temperature of  
20       exhaust air and/or cores.

The spray pressure may vary within wide ranges, in general satisfactory results are obtained with a spray pressure of from about 1 to about 1.5 bar.

The compositions of the invention are useful as immunosuppressants as indicated by standard tests.

The activity and characteristics of the compositions of the invention may be indicated in standard

5 a) clinical trials, e.g. observing the first acute rejection episodes or treatment failure six months after transplant of kidneys or maintaining a rejection - free state within 6 months after initiation of treatment with the invention. The compositions of the invention are administered at a dose in the range of 0.5 to 2.0 g/day e.g. about 1.5 g /day and decrease the acute rejection rates when administered during the period  
10 around transplant surgery, and maintain a rejection-free state in patients who are 3 months or more after transplantation. Thus the compositions of the invention may be administered during the initial 72 hours after transplantation at dose of about 0.5 g administered twice a day in combination with a conventional steroid and cyclosporin, e.g. as NEORAL for which the cyclosporin dose is the conventional  
15 dose e.g. ca.  $8 \pm 3$  mg/kg for renal transplants. The steroid dose is to be administered at about 2.5 mg /kg for 4 days after transplant, 1 mg/kg thereafter for 1 week, 0.6 mg/kg thereafter for 2 weeks thereafter 0.3 mg/kg for 1 month for prednisone.

and in

20 b) animal trials e.g. observing the kidney allograft reaction in rat. In this test one kidney from a female fisher 344 rat is transplanted onto the renal vessel of a unilaterally (left side) nephrectomized WF recipient rat using an end-to-end anastomosis. Ureteric anastomosis is also end-to-end. Treatment commences on the day of transplantation and is continued for 14 days. A contralateral nephrectomy is  
25 done seven days after transplantation, leaving the recipient relying on the performance of the donor kidney. Survival of the graft recipient is taken as the parameter for a functional graft. Typical doses of the compositions of the invention

are from about 1 to 30 mg/kg p.o.

The compositions of the invention are particularly useful for the following conditions:

- a) Treatment and prevention of native or transgenic organ, tissue or cellular allograft or xenograft transplant rejection, e.g. for the treatment of recipients of e.g. heart, lung, combined heart-lung, liver, kidney, pancreatic, skin, pancreatic islet cell, neural cell or corneal transplant; including treatment and prevention of acute rejection; treatment and prevention of hyperacute rejection, e.g. as associated with xenograft rejection; and treatment and prevention of chronic rejection, e.g. as associated with graft-vessel disease. The compositions of the invention are also indicated for the treatment and prevention of graft-versus-host disease, such as following bone marrow transplantation.
- b) Treatment and prevention of autoimmune diseases, e.g. immune-mediated diseases and inflammatory conditions, in particular inflammatory conditions with an etiology including an immunological component such as arthritis (for example rheumatoid arthritis, arthritis chronica progrediente and arthritis deformans) and rheumatic diseases. Specific immune-mediated diseases for which the compositions of the invention may be employed include, autoimmune hematological disorders, including, but not limited to hemolytic anaemia, aplastic anaemia, pure red cell anaemia and idiopathic thrombocytopenia), systemic lupus erythematosus, polychondritis, scleroderma, Wegener granulosis, dermatomyositis, polymyositis, chronic active hepatitis, primary biliary cirrhosis, myasthenia gravis, psoriasis, Steven-Johnson syndrome, pemphigus, idiopathic sprue, inflammatory bowel diseases (including e.g. ulcerative colitis and Crohn's disease), endocrine ophthalmopathy, Graves disease, sarcoidosis, multiple sclerosis, juvenile diabetes (diabetes mellitus type I), non-infectious uveitis (anterior and posterior), keratoconjunctivitis sicca and vernal keratoconjunctivitis, interstitial lung fibrosis, psoriatic arthritis, vasculitis, glomerulonephritides (with and without nephrotic syndrome, e.g. including idiopathic nephrotic syndrome or minimal change nephropathy) and juvenile dermatomyositis.

Appropriate dosages of the compositions of the invention will of course vary, e.g. depending on the condition to be treated (for example the disease type or the nature of resistance), the MPA salt used, the effect desired and the mode of administration.

5 In general however satisfactory results are obtained on administration e.g. orally at dosages on the order of from about 1 to about 30 mg salt per kg animal body weight per day, administered once or in divided doses up to 4 times per day. Suitable daily dosages for patients are thus in the order of 200 mg to 3 g p.o. salt e.g. from about 50 to 100% that of mycophenolate mofetil. For the preferred mono sodium salt the dosage of the salt is about two thirds that of mycophenolate mofetil.

10 Representative unit dosage forms contain from about 50 mg, e.g. 100 mg, to about 1.5 g of the pharmaceutically acceptable mycophenolate salt.

The bioavailability characteristics of compositions of the invention may be determined in conventional manner, e.g. by oral administration to beagle dogs. Dosages are typically 50 mg salt animal e.g. ca 3- 5 mg salt /kg animal body weight. Dogs are adult (ca. 10 kg e.g. 15 6 - 14 kg) and fasted. Three hours after administration ca. 200 g food is administered. Blood samples are taken from the cephalic vein, before administration and 10, 30, and 45 minutes, 1, 1.5, 2, 3, 4, 6, 8, 12, and 24 hours, after administration. Plasma levels of free MPA are determined by HPLC analysis (with UV detection).

20 In a relative bioavailability trial as described above in male beagle dogs dosages of 3.8 mg salt/kg animal body weight p.o. were administered with the Example 1 composition as described hereinafter and with a MPA or MMF formulation corresponding to the Example 1 composition but containing an identical amount of MPA or commercially available MMF.

Results are as follows:-

	MPA (AUC Relative Bioavailability, Frel [ng.hr.ml <sup>-1</sup> ])	Ex 1	MPA	MMF
5	Mean	4612 (218)	3579 (174)	2709 (100)
	Median	4204 (168)	2911 (182)	2513 (100)
	SD	939	1889	1363
	CV	20	53	50
	Cmax [ng/ml] (Relative Cmax)			
10	Mean	5391 (313)	3683 (227)	2052 (100)
	Median	5359 (367)	2719 (172)	1462 (100)
	SD	1847	2504	945
	CV (%)	34 (46)	68 (87)	46 (0)

15 The coefficients of variation (CV) of AUC (20%) and Cmax (34%) of the Example 1 composition are significantly less than those of the reference compositions, indicating less inter-subject and intra-subject variability with the Example 1 composition.

The area under the curve (AUC) and Cmax with the Example 1 composition are higher than those of the reference compositions.

20 Naturally the advantageous bioavailability characteristics of the present compositions may be ascertained in standard clinical bioavailability trials. For example, doses from 200 mg to 1.5 g of the Example 1 composition and MPA, and MMF may be administered to 12 healthy volunteers in single doses in a cross-over trial. Increased AUC and C<sub>max</sub> may be observed for the Example 1 composition.

25 The compositions of the present invention are surprisingly tolerated better than MMF, inducing less gastro-intestinal side effects such as diarrhoea and burning. They show less

long term side effects e.g. in the colon.

The compositions of the invention may be administered as the sole active ingredient or with another immunosuppressant e.g. together with simultaneous or separate administration of other immunosuppressants, for example, in immunosuppressive applications such as prevention and treatment of graft vs. host disease, transplant rejection, or immune-mediated disease, the compositions of the invention may be used in combination with cyclosporins or ascomycins, or their immunosuppressive analogs, e.g. cyclosporin A, FK- 506 (tacrolimus), etc., rapamycin; corticosteroids; cyclophosphamide; azathioprine; methotrexate; brequinar; leflunomide; mizoribine; deoxyspergualin; analogues thereof, and immunosuppressive monoclonal antibodies, e.g., monoclonal antibodies to leukocyte receptors, e.g. MHC, CD2, CD3, CD4, CD7, CD25, CD28, CTLA4, B7, CD45, or CS58 or their ligands; or other immunomodulatory compounds.

When the compositions of the invention are co-administered with such other immunosuppressants the dosages of the other immunosuppressants may be reduced e.g. to one-half to one-third their dosages when used alone.

Representative doses for ciclosporin to be used are e.g. 1 to 10, e.g. 1 to 2 mg/kg/day.

The present invention provides in another aspect the use, method and compositions as defined hereinafter in the claims.

Insofar as details of excipients are not described herein, these are known, or available e.g. in the Handbook of Pharmaceutical Excipients, Second Edition, edited by Ainley Wade and Paul J. Weller, American Pharmaceutical Association, Washington, USA and Pharmaceutical Press, London; and Lexikon der Hilfsstoffe für Pharmazie, Kosmetik and angrenzende Gebiete edited by H.P. Fiedler, 4th Edition, Editio Cantor, Aulendorf and earlier editions.

Following is a description by way of example only of compositions of this invention:

EXAMPLE 1:COMPOSITIONCapsule contents

MPA mono sodium salt 53.43 mg (= 50 mg MPA)

5 Lactose (1:1 mixture of 100/200 mesh) 256.57 mg

Silica (Aerosil) 3.1 mg

Magnesium stearate 1.55 mg

314.65 mg

Capsule is size 1

10 Enteric coating (ca 17 mg)

Hydroxypropyl methyl cellulose phthalate (HP50) 9 parts

Triacetin 1 part

Procedure

15 The capsule ingredients are mixed and filled into size 1 capsules. The capsules are coated in a fluidized bed coater with a solution of the enteric coating ingredients in ethanol (containing 10% acetone). The coating on each capsule is about 17 mg. The capsules meet the enteric coating test described herein and do not disintegrate within 2 hours in artificial gastric juices (pH 1, HCl). The compositions are stable, e.g for 2 years at room temperature.

20 If desired larger capsules containing 534.3 mg MPA mono sodium salt may be made in analogous manner, reducing the amount of lactose. These are well tolerated in clinical trials.

EXAMPLE 2:

Capsules of size 1 are made up as in Example 1. A solution for enteric coating is made up as follows:

Hydroxypropyl methyl cellulose phthalate (HP50) 270 g

25 Triacetin 30 g

Acetone 900 g

Ethanol 1800 g

600 g of this enteric coating solution are used for 1 kg of capsules (ca. 2400). The amount of coating applied to each capsule is about 25 mg giving a film thickness of 5-6 mg/cm<sup>2</sup>.

CLAIMS

1. A pharmaceutical composition comprising a mycophenolate salt, the composition being adapted to release mycophenolate in the upper part of the intestinal tract.
2. A pharmaceutical composition comprising an enteric coated pharmaceutically acceptable mycophenolate salt.
3. Use of an enteric coated pharmaceutically acceptable mycophenolate salt or composition of claim 1 or 2 in the preparation of a medicament for immunosuppression, particularly for prevention or treatment of native or transgenic organ, tissue or cellular allograft or xenograft transplant rejection; for the treatment or prevention of immune-mediated and/or inflammatory diseases; optionally with the simultaneous or separate administration of another immunosuppressant.
4. A method of immunosuppressing a subject which comprises administering a therapeutically effective amount of enteric coated pharmaceutically acceptable mycophenolate or a composition of claim 1 or 2 to a subject in need of such immunosuppression, optionally with the simultaneous or separate administration of another immunosuppressant.
5. A composition containing an enteric coated pharmaceutically acceptable mycophenolate salt or a composition of claim 1 or 2 and another immunosuppressant for simultaneous, sequential or separate administration.
6. A composition, use or method according to any preceding claim wherein the salt is the mono-sodium salt.
7. A composition, use or method according to any preceding claim wherein another immunosuppressant is present, e.g. ciclosporin.